Oliver Buchmueller, Imperial College London

SEARCHES FOR DARK MATTER PRODUCTION AT THE LHC: WHERE ARE WE TODAY?

CPAD INSTRUMENTATION FRONTIER MEETING OCTOBER 5, 2015





Searches for Dark Matter (&SUSY)



Searches for Dark Matter (&SUSY)

Characterizing Dark Matter Searches

complete theory vs. simple interpretations

SUSY



Example:
Effective Field Theory
Simplified models



Characterizing Dark Matter Searches

complete theory vs. simple interpretations

SUSY



Example:
Effective Field Theory
Simplified models



Supersymmetry

Extension of the Standard Model: Introduce a new symmetry

Spin ½ matter particles (fermions) ⇔ Spin 1 force carriers (bosons)

Standard Model particles



SUSY particles



New Quantum number: R-parity: $R_p = (-1)^{B+L+2s}$ = +1 SM particles
- 1 SUSY particles

<u>R-parity conservation</u>:

- SUSY particles are produced in pairs
- The lightest SUSY particle (LSP) is stable

What do we call a "SUSY search"?

The definition is purely derived from the experimental signature.
Therefore, a "SUSY search signature" is characterized by

Lots of missing energy, many jets, and possibly leptons in the final state

Slides from 2007



Missing Energy:
- *from LSP*

Multi-Jet:
- *from cascade decay (gaugino)*

Multi-Leptons:
- *from decay of charginos/neutralios*

RP-Conserving SUSY is a very prominent example predicting this famous signature but …

What is its experimental signature?

… by no means is it the only New Physics model predicting this experimental pattern. Many other NP models predict this genuine signature

Slides from 2007



Missing Energy:

- *Nwimp - end of the cascade*

Multi-Jet:

- *from decay of the Ns (possibly via heavy SM particles like top, W/Z)*

Multi-Leptons:

- *from decay of the N's*

Model examples are Extra dimensions, Little Higgs, Technicolour, etc but a more generic definition for this signature is as follows.

Inclusive SUSY Searches in 2013



Inclusive SUSY Searches in 2013



LEP & Tevatron



The LHC has pushed the mass scale in constraint SUSY models to a new level!

Inclusive SUSY Searches in 2013



Msq = 1800 GeV

Bottom line today:
Impressive variety of powerful SUSY searches have been executed but only limits (at least so far).

LEP & Tevatron



The LHC has pushed the mass scale in constraint SUSY models to a new level!

CMSSM: Evolution with time



X^2 increase from bluish to reddish

Source: http://mastercode.web.cern.ch/mastercode/

Observable	Source Th./Ex.	Constraint	$\Delta\chi^2$ (CMSSM)	$\Delta\chi^2$ (NUHM1)	$\Delta\chi^2$ ("SM")
m_t [GeV]	43	173.2 ± 0.90	0.05	0.06	-
$\Delta\alpha_{had}^{(5)}(M_Z)$	42	0.02749 ± 0.00010	0.009	0.004	-
M_Z [GeV]	44	91.1875 ± 0.0021	2.7×10^{-5}	0.26	-
Γ_Z [GeV]	26 / 44	$2.4952 \pm 0.0023 \pm 0.001_{SUSY}$	0.078	0.047	0.14
σ_{had}^0 [nb]	26 / 44	41.540 ± 0.037	2.50	2.57	2.54
R_l	26 / 44	20.767 ± 0.025	1.05	1.08	1.08
$A_{fb}(\ell)$	26 / 44	0.01714 ± 0.00095	0.72	0.69	0.81
$A_\ell(P_\tau)$	26 / 44	0.1465 ± 0.0032	0.11	0.13	0.07
R_b	26 / 44	0.21629 ± 0.00066	0.26	0.29	0.27
R_c	26 / 44	0.1721 ± 0.0030	0.002	0.002	0.002
$A_{fb}(b)$	26 / 44	0.0992 ± 0.0016	7.17	7.37	6.63
$A_{fb}(c)$	26 / 44	0.0707 ± 0.0035	0.86	0.88	0.80
A_b	26 / 44	0.923 ± 0.020	0.36	0.36	0.35
A_c	26 / 44	0.670 ± 0.027	0.005	0.005	0.005
A_ℓ(SLD)	26 / 44	0.1513 ± 0.0021	3.16	3.03	3.51
$\sin^2\theta_w^\ell(Q_{fb})$	26 / 44	0.2324 ± 0.0012	0.63	0.64	0.59
M_W [GeV]	26 / 44	$80.399 \pm 0.023 \pm 0.010_{SUSY}$	1.77	1.39	2.08
$a_\mu^{EXP} - a_\mu^{SM}$	53 / 42,54	$(30.2 \pm 8.8 \pm 2.0_{SUSY}) \times 10^{-10}$	4.35	1.82	11.19 (N/A)
M_h [GeV]	28 / 55,56	$> 114.4[\pm1.5_{SUSY}]$	0.0	0.0	0.0
$BR_{b\to s\gamma}^{EXP/SM}$	45 / 46	$1.117 \pm 0.076_{EXP} \pm0.082_{SM} \pm 0.050_{SUSY}$	1.83	1.09	0.94
$BR(B_s \to \mu^+\mu^-)$	29 / 41	CMS & LHCb	0.04	0.44	0.01
$BR_{B\to\tau\nu}^{EXP/SM}$	29 / 46	$1.43 \pm 0.43_{EXP+TH}$	1.43	1.59	1.00
$BR(B_d \to \mu^+\mu^-)$	29 / 46	$< 4.6[\pm0.01_{SUSY}] \times 10^{-9}$	0.0	0.0	0.0
$BR_{B\to X_s\ell\ell}^{EXP/SM}$	47 / 46	0.99 ± 0.32	0.02	$\ll 0.01$	$\ll 0.01$
$BR_{K\to\mu\nu}^{EXP/SM}$	29 / 48	$1.008 \pm 0.014_{EXP+TH}$	0.39	0.42	0.33
$BR_{K\to\pi\nu\nu}^{EXP/SM}$	49 / 50	< 4.5	0.0	0.0	0.0
$\Delta M_{B_s}^{EXP/SM}$	49 / 51,52	$0.97 \pm 0.01_{EXP} \pm 0.27_{SM}$	0.02	0.02	0.01
$\frac{\Delta M_{B_s}^{EXP/SM}}{\Delta M_{B_d}^{EXP/SM}}$	29 / 46,51,52	$1.00 \pm 0.01_{EXP} \pm 0.13_{SM}$	$\ll 0.01$	0.33	$\ll 0.01$
$\Delta\epsilon_K^{EXP/SM}$	49 / 51,52	$1.08 \pm 0.14_{EXP+TH}$	0.27	0.37	0.33
$\Omega_{CDM}h^2$	31 / 13	$0.1120 \pm 0.0056 \pm 0.012_{SUSY}$	8.4×10^{-4}	0.1	N/A
σ_p^{SI}	25	$(m_{\tilde\chi^0_1}, \sigma_p^{SI})$ plane	0.13	0.13	N/A
jets + $\not{E}_T$	18,20	$(m_0, m_{1/2})$ plane	1.55	2.20	N/A
$H/A, H^\pm$	21	$(M_A, \tan\beta)$ plane	0.0	0.0	N/A
Total χ^2/d.o.f. p-values	All	All	28.8/22 15%	27.3/21 16%	32.7/23 (21.5/22) 9% (49%)

Global Fit to indirect and direct constraints on SUSY!

Other "fitter" groups find very similar results: e.g.
SuperBayeS: **arXiv:1212.2636**
Fittino group: **arXiv:1204.4199**

CMSSM: Evolution with time



X^2 increase from bluish to reddish

O. Buchmueller[a], R. Cavanaugh[b,c], A. De Roeck[d,e], M.J. Dolan[f], J.R. Ellis[g,d], H. Flächer[h], S. Heinemeyer[i], G. Isidori[j,d], J. Marrouche[a], D. Martínez Santos[k], K.A. Olive[l], S. Rogerson[a], F.J. Ronga[m], K.J. de Vries[a], G. Weiglein[n]

[a]High Energy Physics Group, Blackett Laboratory, Imperial College, Prince Consort Road, London SW7 2AZ, UK

[b]Fermi National Accelerator Laboratory, P.O. Box 500, Batavia, Illinois 60510, USA

[c]Physics Department, University of Illinois at Chicago, Chicago, Illinois 60607-7059, USA

[d]Physics Department, CERN, CH–1211 Genève 23, Switzerland

[e]Antwerp University, B–2610 Wilrijk, Belgium

[f]Theory Group, SLAC National Accelerator Laboratory, 2575 Sand Hill Road, Menlo Park, CA 94025-7090, USA

[g]Theoretical Particle Physics and Cosmology Group, Department of Physics, King's College London, London WC2R 2LS, UK

[h]H.H. Wills Physics Laboratory, University of Bristol, Tyndall Avenue, Bristol BS8 1TL, UK

[i]Instituto de Física de Cantabria (CSIC-UC), E–39005 Santander, Spain

[j]INFN, Laboratori Nazionali di Frascati, Via E. Fermi 40, I–00044 Frascati, Italy

[k]NIKHEF and VU University Amsterdam, Science Park 105, NL-1098 XG Amsterdam, The Netherlands

[l]William I. Fine Theoretical Physics Institute, School of Physics and Astronomy, University of Minnesota, Minneapolis, Minnesota 55455, USA

[m]Institute for Particle Physics, ETH Zürich, CH–8093 Zürich, Switzerland

[n]DESY, Notkestrasse 85, D–22607 Hamburg, Germany

Global Fit to indirect ar
constraints on SUSY!

Other "fitter" groups find
results: e.g.
SuperBayeS: **arXiv:121**
Fittino group: **arXiv:120**

CMSSM: Evolution with time



CMSSM today


stau coann.
A/H-funnel
hybrid
$\tilde{\chi}_1^{\pm}$ coann.
focus point
LUX/XENON100 90% CL excluded
neutrino floor
1310.8327
σ_p^{SI} [cm^2]
$m_{\tilde{\chi}_1^0}$ [GeV]
10^{-39}
10^{-40}
10^{-41}
10^{-42}
10^{-43}
10^{-44}
10^{-45}
10^{-46}
10^{-47}
10^{-48}
10^{-49}
10^{-50}
10^0
10^1
10^2
10^3
10^4

NUHM1 today


stau coann.
A/H-funnel
hybrid
$\tilde{\chi}_1^{\pm}$ coann.
focus point
LUX/XENON100 90% CL excluded
neutrino floor
1310.8327
σ_p^{SI} [cm^2]
$m_{\tilde{\chi}_1^0}$ [GeV]
10^{-39}
10^{-40}
10^{-41}
10^{-42}
10^{-43}
10^{-44}
10^{-45}
10^{-46}
10^{-47}
10^{-48}
10^{-49}
10^{-50}
10^0
10^1
10^2
10^3
10^4

NUHM2 today


stau coann.
A/H-funnel
hybrid
$\tilde{\chi}_1^{\pm}$ coann.
focus point
LUX/XENON100 90% CL excluded
neutrino floor
1310.8327
σ_p^{SI} [cm^2]
$m_{\tilde{\chi}_1^0}$ [GeV]
10^{-39}
10^{-40}
10^{-41}
10^{-42}
10^{-43}
10^{-44}
10^{-45}
10^{-46}
10^{-47}
10^{-48}
10^{-49}
10^{-50}
10^{0}
10^{1}
10^{2}
10^{3}
10^{4}

pMSSM10 today



SUSY Status – post 7 TeV LHC data

- Constrained SUSY models like the CMSSM are severely put under pressure by the LHC limits!
- Experiments need to define new benchmarks and to present the interpretation of their searches.
- A bottom-up approach, using so-called simplified models, was adopted but ATLAS and CMS as the primary vehicle to present SUSY searches!

Interpretation in Simplified Models

DM Searches @ LHC O. Buchmüller

CMSSM

What the individual searches are sensitive to is much more simple…



86% of all hadronic production in LM1 consists of "simple" decay chains.

This makes it particularly amenable to being ***approximated well*** with a 3-particle OSET.

Simplified model spectrum (SMS)
with 3 particles, 2 decay modes

SMS: a few interesting features

$m_{LSP}^{max} \approx 0.3$ TeV : LSP mass above which there is NO limit anymore





$\tilde{q}\tilde{q} \rightarrow q\tilde{\chi}^0 \bar{q}\tilde{\chi}^0$

$m_G^{max} \approx 0.8$ TeV : Best limit in plane

Assumes 100% BR for decay chain considered.

How to summarize SMS limits?

Approach taken in the 2012 and 2013 Experimental SUSY PDG reviews [OB & Paul De Jong]:

http://pdg.lbl.gov/2012/reviews/rpp2012-rev-susy-2-experiment.pdf
http://pdg.lbl.gov/2013/reviews/rpp2013-rev-susy-2-experiment.pdf

Model	Assumption	$m_{\tilde{q}}$	$m_{\tilde{g}}$
CMSSM	$m_{\tilde{q}} \approx m_{\tilde{g}}$	1400	1400
	all $m_{\tilde{q}}$	-	800
	all $m_{\tilde{g}}$	1300	-
Simplified model $\tilde{g}\tilde{g}$	$m_{\tilde{\chi}_1^0}$ =0	-	900
	$m_{\tilde{\chi}_1^0} > 300$	-	no limit
Simplified model $\tilde{q}\tilde{q}$	$m_{\tilde{\chi}_1^0} = 0$	750	-
	$m_{\tilde{\chi}_1^0} > 250$	no limit	-
Simplified model $\tilde{g}\tilde{q}, \tilde{g}\bar{\tilde{q}}$	$m_{\tilde{\chi}_1^0}$= 0, $m_{\tilde{q}} \approx m_{\tilde{g}}$	1500	1500
	$m_{\tilde{\chi}_1^0}$= 0, all $m_{\tilde{g}}$	1400	-
	$m_{\tilde{\chi}_1^0}$= 0, all $m_{\tilde{q}}$	-	900



This was an appropriate approach for the rather limited amount of inclusive searches and corresponding SMS interpretations available in 2011 (7 TeV).

How to summarize SMS limits?

Approach taken in the 2012 and 2013 Experimental SUSY PDG reviews [OB & Paul De Jong]:

http://pdg.lbl.gov/2012/reviews/rpp2012-rev-susy-2-experiment.pdf
http://pdg.lbl.gov/2013/reviews/rpp2013-rev-susy-2-experiment.pdf

Model	Assumption	$m_{\tilde{q}}$	$m_{\tilde{g}}$
	$m_{\tilde{q}} \approx m_{\tilde{g}}$	1400	1400
CMSSM	all m_0		800
Simpli			
Simplified model $\tilde{q}\tilde{q}$	$m_{\tilde{\chi}_1^0} = 0$	750	-
	$m_{\tilde{\chi}_1^0} > 250$	no limit	-
Simplified model $\tilde{g}\tilde{q}, \tilde{g}\bar{\tilde{q}}$	$m_{\tilde{\chi}_1^0} = 0$, $m_{\tilde{q}} \approx m_{\tilde{g}}$	1500	1500
	$m_{\tilde{\chi}_1^0} = 0$, all $m_{\tilde{g}}$	1400	-
	$m_{\tilde{\chi}_1^0} = 0$, all $m_{\tilde{q}}$	-	900

It is a challenge to do justice to the many searches and limits that have been established so far
- even more so to put it all together into the/a "bigger picture".



This was an appropriate approach for the rather limited amount of inclusive searches and corresponding SMS interpretations available in 2011 (7 TeV).

Note: The following results are a **May 2015 update** to PDG review September 2013.

http://pdg.lbl.gov/2013/reviews/rpp2013-rev-susy-2-experiment.pdf



What are representative SMS limits on the different particles?


SUSY & DM Searches @ LHC O. Buchmüller
m_{LSP} [GeV]
Direct squark
$m_{SUSY} = m_{\tilde{q}}$
$\tilde{q} \to q\chi_1^0$
CMS arXiv:1502.04358
$\tilde{u}_L \to u\chi_1^0$
CMS arXiv:1502.04358
$\tilde{b} \to b\chi_1^0$
ATLAS arXiv:1308.2631
$\tilde{t} \to t\chi_1^0$
ATLAS arXiv:1407.0583
Direct stop in "gap"
$m_{SUSY} = m_{\tilde{t}}$
$\tilde{t} \to c\chi_1^0$
ATLAS arXiv:1407.0608
$\tilde{t} \to bff'\chi_1^0$
ATLAS arXiv:1407.0583
$\tilde{t} \to Wb\chi_1^0$
CMS arXiv:1308.1586
Gluino mediated
$m_{SUSY} = m_{\tilde{g}}$
$\tilde{g} \to q\bar{q}\chi_1^0$
ATLAS arXiv:1405.7875
$\tilde{g} \to b\bar{b}\chi_1^0$
CMS arXiv:1502.00300
$\tilde{g} \to t\bar{t}\chi_1^0$
ATLAS arXiv:1407.0600
Direct slepton
$m_{SUSY} = m_{\tilde{l}}$
$\tilde{l}_R \to l^{\pm}\chi_1^0$
ATLAS: arXiv:1403.5294
$\tilde{l}_L \to l^{\pm}\chi_1^0$
ATLAS: arXiv:1403.5294
$m_{SUSY} - m_{LSP} < m_t$
$m_{SUSY} - m_{LSP} < 2m_t$
$m_{\tilde{t}} - m_b - m_{LSP} < m_W$
$m_{\tilde{t}} - m_{LSP} < m_t$
Direct $\chi_1^+\chi_1^-$
$m_{SUSY} = m_{\chi_1^{\pm}}$
$\chi_1^+\chi_1^-$(heavy $\tilde{l}$)
ATLAS arXiv:1403.5294
$\chi_1^+\chi_1^-$(light $\tilde{l}$)
ATLAS arXiv:1403.5294
BR=100%
all limits are observed nominal 95% CLs limits
RP conserved
Direct $\chi_1^{\pm}/\chi_2^0$
$m_{SUSY} = m_{\chi_1^{\pm}} = m_{\chi_2^0}$
$\chi_1^{\pm}\chi_2^0$(light $\tilde{l}$)
CMS: arXiv:1405.7570
$\chi_1^{\pm}\chi_2^0$(heavy $\tilde{l}$)
CMS: arXiv:1405.7570
m_{SUSY} [GeV]
0
250
500
750
1000
1250
1500


SUSY & DM Searches @ LHC O. Buchmüller
m_{LSP} [GeV]
m_{SUSY} [GeV]
Direct squark
$m_{SUSY} = m_{\tilde{q}}$
$\tilde{q} \to q\chi_1^0$
CMS arXiv:1502.04358
$\tilde{u}_L \to u\chi_1^0$
CMS arXiv:1502.04358
$\tilde{b} \to b\chi_1^0$
ATLAS arXiv:1308.2631
$\tilde{t} \to t\chi_1^0$
ATLAS arXiv:1407.0583
Direct stop in "gap"
$m_{SUSY} = m_{\tilde{t}}$
$\tilde{t} \to c\chi_1^0$
ATLAS arXiv:1407.0608
$\tilde{t} \to bff'\chi_1^0$
ATLAS arXiv:1407.0583
$\tilde{t} \to Wb\chi_1^0$
CMS arXiv:1308.1586
Direct slepton
$m_{SUSY} = m_{\tilde{l}}$
$\tilde{l}_R \to l^{\pm}\chi_1^0$
ATLAS: arXiv:1403.5294
$\tilde{l}_L \to l^{\pm}\chi_1^0$
ATLAS: arXiv:1403.5294
$m_{\tilde{t}} - m_b - m_{LSP} < m_W$
$m_{\tilde{t}} - m_{LSP} < m_t$
$m_{SUSY} - m_{\tilde{t}}$
$m_{SUSY} - m_{\tilde{L}}$
$\chi_1^+\chi_1^-$ (heavy $\tilde{l}$)
ATLAS arXiv:1403.5294
$\chi_1^+\chi_1^-$ (light $\tilde{l}$)
ATLAS arXiv:1403.5294
BR=100%
all limits are observed nominal 95% CLs limits RP conserved
Direct $\chi_1^{\pm}/\chi_2^0$
$m_{SUSY} = m_{\chi_1^{\pm}} = m_{\chi_2^0}$
$\chi_1^{\pm}\chi_2^0$(light $\tilde{l}$)
CMS: arXiv:1405.7570
$\chi_1^{\pm}\chi_2^0$(heavy $\tilde{l}$)
CMS: arXiv:1405.7570
MSUGRA/CMSSM: tan(β) = 30, A_0 = -2m_0, μ > 0
Lepton & Photon 2013
ATLAS Preliminary
$\int L\,dt$ = 20.1 - 20.7 fb^{-1}, $\sqrt{s}$ = 8 TeV
95% CL limits. σ^{SUSY}_{theory} not included.
Expected / Observed 0-lepton, 2-6 jets ATLAS-CONF-2013-047
Expected / Observed 0-lepton, 7-10 jets ATLAS-CONF-2013-054
Expected / Observed 0-1 lepton, 3 b-jets ATLAS-CONF-2013-061
Expected / Observed 1-lepton +jets +MET ATLAS-CONF-2013-062
Expected / Observed 1-2 taus +jets +MET ATLAS-CONF-2013-026
Expected / Observed 2-SS-leptons, 0 - ≥3 b-jets ATLAS-CONF-2013-007
$m_{1/2}$ [GeV]
m_0 [GeV]

What does this imply for Linear Collider?




SUSY & DM Searches @ LHC O. Buchmüller
m_{LSP} [GeV]
Direct squark
$m_{SUSY} = m_{\tilde{q}}$
$\tilde{q} \to q\chi_1^0$
CMS arXiv:1502.04358
$\tilde{u}_L \to u\chi_1^0$
CMS arXiv:1502.04358
$\tilde{b} \to b\chi_1^0$
ATLAS arXiv:1308.2631
$\tilde{t} \to t\chi_1^0$
ATLAS arXiv:1407.0583
Direct stop in "gap"
$m_{SUSY} = m_{\tilde{t}}$
$\tilde{t} \to c\chi_1^0$
ATLAS arXiv:1407.0608
$\tilde{t} \to bff'\chi_1^0$
ATLAS arXiv:1407.0583
$\tilde{t} \to Wb\chi_1^0$
CMS arXiv:1308.1586
Gluino mediated
$m_{SUSY} = m_{\tilde{g}}$
$\tilde{g} \to q\bar{q}\chi_1^0$
ATLAS arXiv:1405.7875
$\tilde{g} \to b\bar{b}\chi_1^0$
CMS arXiv:1502.00300
$\tilde{g} \to t\bar{t}\chi_1^0$
ATLAS arXiv:1407.0600
Direct slepton
$m_{SUSY} = m_{\tilde{l}}$
$\tilde{l}_R \to l^{\pm}\chi_1^0$
ATLAS: arXiv:1403.5294
$\tilde{l}_L \to l^{\pm}\chi_1^0$
ATLAS: arXiv:1403.5294
$m_{SUSY} - m_{LSP} < m_t$
$m_{SUSY} - m_{LSP} < 2m_t$
$m_{\tilde{t}} - m_b - m_{LSP} < m_W$
$m_{\tilde{t}} - m_{LSP} < m_t$
Direct $\chi_1^+\chi_1^-$
$m_{SUSY} = m_{\chi_1^{\pm}}$
$\chi_1^+\chi_1^-$(heavy $\tilde{l}$)
ATLAS arXiv:1403.5294
$\chi_1^+\chi_1^-$(light $\tilde{l}$)
ATLAS arXiv:1403.5294
BR=100%
all limits are observed nominal 95% CLs limits RP conserved
Direct $\chi_1^{\pm}/\chi_2^0$
$m_{SUSY} = m_{\chi_1^{\pm}} = m_{\chi_2^0}$
$\chi_1^{\pm}\chi_2^0$(light $\tilde{l}$)
CMS: arXiv:1405.7570
$\chi_1^{\pm}\chi_2^0$(heavy $\tilde{l}$)
CMS: arXiv:1405.7570
m_{SUSY} [GeV]
0
250
500
750
1000
1250
1500



There remain kinematic regions that are currently beyond the reach of the LHC while still being accessible with a 1 TeV linear collider.

It will take an effort (and low trigger thresholds) to fill as most as possible of it in the future!

Characterizing Dark Matter Searches

complete theory vs. simple interpretations

SUSY



Example:
Effective Field Theory
Simplified models



Mono-X searches at colliders



E_T^{miss} trigger

Example Monojet (8 TeV, 20.3 fb-1)

- E_T^{miss} , $p_T(j) > 150 - 900$ GeV
- 1 or 2 jets (anti-k_T, R=0.4, p_T>30 GeV)
- $|\Delta\phi(E_T^{miss}, j_2)|>0.5$

Example Monophoton (8 TeV, 19.6 fb^{-1}):

- E_T^{miss} , $p_T(\gamma)>140$ GeV,
- $N_{jet} < 2$ (anti-k_T, R=0.5, p_T>30 GeV)
- $\Delta\phi (\gamma, E_T^{miss}) >2$,
- (X^2, $\Delta\phi$ (jet, E_T^{miss}) >0.4)

ATLAS Mono-Jet: Comparison with Direct Detection



ATLAS Mono-Jet: Comparison with Direct Detection



ATLAS Mono-Jet: Comparison with Direct Detection



Mono-Jet analyses better than direct detection?!





Claim [often made]:

For low mass and the entire spin-dependent case monojet limits are stronger than direct detection limits!

Validity of Effective Field Theory Limits

Recent work from OB, M.Dolan,C.McCabe: arXiv:1308.6799

- Compare Effective Field Theory (EFT) with Full Theory (FT)

EFT approach





FT
one diagram
"simplified model"

Use vector and axial-vector mediators (e.g. Z') as example - scalar are similar in conclusion!



Three Regions as function of mediator mass:

Region I: Heavy m_{med}

- EFT is valid!

Region II: Medium m_{med} – Resonant enhancement

- EFT limits are too conservative!

Region III: Low m_{med}

- EFT limits are far too aggressive!

Validity of Effective Field Theory Limits

Recent work from OB, M.Dolan,C.McCabe: arXiv:1308.6799

- Compare Effective Field Theory (EFT) with Full Theory (FT)

EFT approach





FT
one diagram
"simplified model"

Use v … clusion!

Conclusion:
The EFT is not an appropriate framework for a comprehensive Interpretation of DM searches at colliders and especially must taken with very (as in VERY) special care when comparing with other experiments such as Direct Detection!



Region I: Heavy m_{med}
- EFT is valid!

Region II: Medium m_{med} – Resonant enhancement
- EFT limits are too conservative!

Region III: Low m_{med}
- EFT limits are far too aggressive!

Minimal Simplified Dark Matter Model

Based on work from :
OB, S. Malik, M.Dolan,C.McCabe arXiV:1407.8257



s-channel

Define simplified model with (minimum) 4 parameters	
Mediator mass (M_{med})	DM mass (M_{DM})
g_q	g_{DM}

DM	
Dirac fermion	Scalar - real
Majorana fermion	Scalar - complex

Consider comprehensive set of diagrams for mediator	
Vector	Axial-vector
Scalar	Pseudoscalar

(Γ_{med} can also be free as long As $\Gamma_{med}<M_{med}$)

Minimal Simplified Dark Matter Model

Based on work from :
OB, S. Malik,
M.Dolan,C.McCabe
arXiV:1407.8257



s-channel

All this is also discussed in the recent DM forum, which brought together experimentalist from ATLAS and CMS as well as theorists. See: arXiv:1507.00966

Def (minimum) 4 parameters		DM		sive set of diagrams for mediator	
Mediator mass (M_{med})	DM mass (M_{DM})	Dirac fermion	Scalar - real	Vector	Axial-vector
g_q	g_{DM}	Majorana fermion	Scalar - complex	Scalar	Pseudoscalar

(Γ_{med} can also be free as long As $\Gamma_{med}<M_{med}$)

Collider vs Direct Detection

M_{DM}	M_{med}
g_q	g_{DM}

Vector



Axial vector



arXiV:1407.8257

Scalar and Pseudoscalar

Philip Harris, Valentin V. Khoze,
Michael Spannowsky, Ciaran Williams

arXiv:1411.0535

See also Buckley et al
arXiv:1410.6497

chmüller

Scalar



Pseudoscalar



Projections for Future Experiments: M_{med} vs M_{DM}

Based on work from :
S. Malik, OB, M.Dolan, C.McCabe et al
arXiV:1409.4075

Limits from 8 TeV monojet search and projected limits for 3 LHC scenarios:
- 13 TeV 30 fb^{-1}
- 14 TeV, 300 fb^{-1}
- 14 TeV, 3000 fb^{-1}

LUX 2013 limits and projected limits for LZ assuming 10 tonne-year exposure

Discovery reach accounting for coherent neutrino scattering



Projections for Future Experiments: σ vs M_{DM}

Can be also shown in the σ vs M_{DM} plane …





Direct Detection experiments and collider are complementary!
They are probing different regions of the relevant parameter space!

Summary for most basic Mediator Interactions

… in a nutshell!

Basic Mediators	
Vector EWK like coupling (assumed equal to all leptons). *Besides very low DM masses DD wins clearly over collider!*	**Axial-vector** EWK like coupling (assumed equal to all leptons). *DD and collider are equal in overall sensitivity but probe different regions of parameter space!*
Scalar Yukawa like coupling on SM side (mass based on SM side) *DD and collider are equal in overall sensitivity but probe different regions of parameter space!*	**Pseudoscalar** Yukawa like coupling on SM side (mass based on SM side) *No limits from DD (only from indirect detection). Collider provides limits similar in sensitivity to scalar limits*

Projections for future experiments

Example:
Axial-Vector Mediator
$g_{SM}=g_{DM}=1$




Neutrino background
m_{DM} [GeV]
4000
3000
2000
1000
0
0
2500
5000
7500
10000
12500
15000
m_{MED} [GeV]
SUSY & DM Searches @ LHC O. Buchmüller
WIMP–nucleon cross section [cm^2]
WIMP–nucleon cross section [pb]
WIMP Mass [GeV/c^2]
SuperCDMS Soudan Low Threshold
XENON 10 S2 (2013)
CDMS-II Ge Low Threshold (2011)
CoGeNT (2012)
CDMS Si (2013)
DAMA
SIMPLE (2012)
COUPP (2012)
ZEPLIN-III (2012)
CDMS II Ge (2009)
EDELWEISS (2011)
Xenon100 (2012)
LUX (2013)
DarkSide 50
PICO250-CF3I
SuperCDMS Soudan
LUX 300-day
SuperCDMS SNOLAB
Xenon1T
DarkSide G2
LZ
DEAP3600
CRESST
CDMSlite (2013)
PICO250-C3F8
SuperCDMS SNOLAB
NEUTRINO COHERENT SCATTERING
^{7}Be Neutrinos
^{8}B Neutrinos
Atmospheric and DSNB Neutrinos
(Green ovals) Asymmetric DM
(Violet oval) Magnetic DM
(Blue oval) Extra dimensions
(Red circle) SUSY MSSM
MSSM: Pure Higgsino
MSSM: A funnel
MSSM: Bino-stop coannihilation
MSSM: Bino-squark coannihilation
Axial-Vector Mediator
$g_{SM}=g_{DM}=1$


Neutrino background
LHC 8 TeV 19.5 fb-1
LUX2013
taken from arXiV:1409.4075
Axial-Vector Mediator
gSM=gDM=1
Today
mDM [GeV]
mMED [GeV]
4000
3000
2000
1000
0
0
2500
5000
7500
10000
12500
15000
SUSY & DM Searches @ LHC O. Buchmüller


Neutrino background
LHC 8 TeV 19.5 fb-1
LHC 14 TeV 300 fb-1
LUX2013
LZ 10 ton yr
taken from arXiV:1409.4075
Axial-Vector Mediator
gSM=gDM=1
2020ish
mDM [GeV]
mMED [GeV]
4000
3000
2000
1000
0
0
2500
5000
7500
10000
12500
15000
SUSY & DM Searches @ LHC O. Buchmüller


Neutrino background
LHC 8 TeV 19.5 fb-1
LHC 14 TeV 300 fb-1
LHC 14 TeV 3000 fb-1
LUX2013
LZ 10 ton yr
taken from arXiV:1409.4075
mDM [GeV]
4000
3000
2000
1000
0
0
2500
5000
7500
10000
12500
15000
mMED [GeV]
Axial-Vector Mediator
gSM=gDM=1
2025ish
SUSY & DM Searches @ LHC O. Buchmüller


FCC 100 TeV 1ab-1
taken from arXiV:1509.02904
Neutrino background
LHC 8 TeV 19.5 fb-1
LHC 14 TeV 300 fb-1
LHC 14 TeV 3000 fb-1
LUX2013
LZ 10 ton yr
taken from arXiV:1409.4075
mDM [GeV]
4000
3000
2000
1000
0
0
2500
5000
7500
10000
12500
15000
mMED [GeV]
Axial-Vector Mediator
gSM=gDM=1
Some day perhaps?
SUSY & DM Searches @ LHC O. Buchmüller

Fermi-Lat Excess: Indirect Detection vs Collider

Constraining the Fermi-LAT excess with collider data.

Based on work from :
OB, S. Malik, C.McCabe, and B. Penning
arXiV:1505.07826



$M_{DM} \sim 45$ GeV
arXiv:1409.0042





Fermi-Lat Excess: Indirect Detection vs Collider

Constraining the Fermi-LAT excess with collider data.

Based on work from :
OB, S. Malik, C.McCabe, and B. Penning
arXiV:1505.07826
Paper appeared yesterday on arXiv



$M_{DM} \sim 45$ GeV
arXiv:1409.0042



Region excluded for pseudo-scalar model interpretation:



Outlook: 8 TeV vs 14 TeV

Use parton luminosities to illustrate the gain of 14 vs 8 TeV

Higgs:
pp → H, H→WW, ZZ and γγ
mainly gg: factor ~2

SUSY – 3rd Generation:
Mass scale ~ 500 GeV
qq and gg: factor ~3 to 6

Scalar/Pseudoscalar Mediator
Mass scale ~ 2.0 TeV
gg: factor ~20

SUSY – Squarks/Gluino:
Mass scale ~ 1.5 TeV
qq,gg,qg: factor ~40 to 80

Vector/Axialvector a la Z' :
Mass scale ~ 5 TeV
qq: factor ~1000



Increase in energy will help a lot!

Summary

- So far New Physics has not revealed itself!
  - In 2010 the LHC has enter new territory for New Physics searches and since then pushed e.g. the (coloured) mass scale to the ~1 TeV scale and in some cases even beyond.
- The LHC experiments have established an impressive variety of very powerful direct searches for many different final states!
  - So far many of these searches are only interpret in dedicated models, while they do possess also sensitivity for other scenarios. A good example are the missing (transverse) energy searches, which, so far, predominately have been interpret in the context of "SUSY" only.
  - Based on these results we need to establish the "big picture" in order to understand if/ where our search strategy might have weak spots or even holes!
  - This requires appropriate interpretations of the searches and a MEANIGFUL comparison with other experiments – important example are DM searches!
  - The LHC DM forum has made a first important step in this direction!
- The high energy running of the LHC starting 2015 will be our next very (as in VERY) real chance for discovery!

The story continues … stay tuned!

BACKUP

ATLAS & CMS public results

Most results presented in this talk (and many more) can be accessed via the public page of the ATLAS and CMS experiments:

ATLAS SUSY: https://twiki.cern.ch/twiki/bin/view/AtlasPublic/SupersymmetryPublicResults

CMS SUSY :https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsSUS

100 TeV Prediction from arXiv:1509.02904









The full story

SUSY SUMMARY PLOT

Note: The following results are a **May 2015 update** to PDG review September 2013.

http://pdg.lbl.gov/2013/reviews/rpp2013-rev-susy-2-experiment.pdf



What are representative SMS limits on the different particles?

Direct squark production – chosen limits













CMS arXiv:1502.04358
Signature: Jets + E_t^{miss} with M_{T2}
Limit assumes all 1st & 2nd gen squarks to be mass degenerate [or only one light squark]!

ATLAS arXiv:1308.2631
Signature: 2 b-jets + E_T^{mis}

ATLAS arXiv:1407.0583
Signature: 1Lepton + jets + E_T^{mis}


Direct squark
$m_{SUSY} = m_{\tilde{q}}$
$\tilde{q} \to q\chi_1^0$
CMS arXiv:1502.04358
$\tilde{u_L} \to u\chi_1^0$
CMS arXiv:1502.04358
$\tilde{b} \to b\chi_1^0$
ATLAS arXiv:1308.2631
$\tilde{t} \to t\chi_1^0$
ATLAS arXiv:1407.0583
$m_{SUSY} - m_{LSP} < m_t$
BR=100%
all limits are observed nominal 95% CLs limits
RP conserved
m_{LSP} [GeV]
m_{SUSY} [GeV]
0
250
500
750
1000
1250
1500
SUSY & DM Searches @ LHC O. Buchmüller

Gluino mediated squark production – limits chosen













ATLAS arXiv:1405.7875
Signature: 0L + 2-6 Jets + E_t^{miss}

CMS arXiv:1502.00300
Signature: : 0L + Razor + b-tag

Signature: 0/1 Leptons + 3 b-tag + E_t^{mis}


SUSY & DM Searches @ LHC O. Buchmüller
m_{LSP} [GeV]
Direct squark
$m_{SUSY} = m_{\tilde{q}}$
$\tilde{q} \to q\chi_1^0$
CMS arXiv:1502.04358
$\tilde{u}_L \to u\chi_1^0$
CMS arXiv:1502.04358
$\tilde{b} \to b\chi_1^0$
ATLAS arXiv:1308.2631
$\tilde{t} \to t\chi_1^0$
ATLAS arXiv:1407.0583
Gluino mediated
$m_{SUSY} = m_{\tilde{g}}$
$\tilde{g} \to q\bar{q}\chi_1^0$
ATLAS arXiv:1405.7875
$\tilde{g} \to b\bar{b}\chi_1^0$
CMS arXiv:1502.00300
$\tilde{g} \to t\bar{t}\chi_1^0$
ATLAS arXiv:1407.0600
$m_{SUSY} - m_{LSP} < m_t$
$m_{SUSY} - m_{LSP} < 2m_t$
BR=100%
all limits are observed nominal 95% CLs limits
RP conserved
m_{SUSY} [GeV]
0
250
500
750
1000
1250
1500


Mind the gap!
$\Delta m = m(\tilde{t}) - m(\tilde{\chi}^0)$
$\Delta m < m(W)$
$m(W) < \Delta m < m(t)$
$\Delta m > m(t)$
$\tilde{t} \to c\tilde{\chi}_1^0$
$\tilde{t} \to bW^*\tilde{\chi}_1^0$
$\tilde{t} \to b\tilde{\chi}_1^+ \to bW\tilde{\chi}_1^0$
$\tilde{t} \to bW\tilde{\chi}_1^0$ (off-shell top)
$\tilde{t} \to t\tilde{\chi}_1^0$ (on-shell top)
$\tilde{t}\tilde{t}^* \to b\bar{b}\tilde{\chi}_1^+\tilde{\chi}_1^- \to b\bar{b}W^+W^-\tilde{\chi}_1^0\tilde{\chi}_1^0$
$\tilde{t}\tilde{t}^* \to t\bar{t}\tilde{\chi}_1^0\tilde{\chi}_1^0$
MET
$m(\tilde{\chi}^0)$ [GeV]
$m(\tilde{t})$ [GeV]
0
m(W)
m(t)
m_{LSP} [GeV]
m_{SUSY} [GeV]
1000
750
500
250
0
250
500
750
1000
1250
1500
SUSY & DM Searches @ LHC O. Buchmüller
ediated
$= m_{\tilde{g}}$
1405.7875
02.00300
1407.0600
BR=100%
all limits are observed nominal 95% CLs limits
RP conserved
64

Compressed stop – mind the gap!



$\tilde{t} \to c\chi_1^0$



$\tilde{t} \to bff'\chi_1^0$



$\tilde{t} \to Wb\chi_1^0$







ATLAS arXiv:1407.0608
Mono-jet & c-tag combined

ATLAS: arXiv:1407.0583
1L + E_t^{mis} & b-tag

CMS arXiv:1308.1586
1L + E_t^{mis} and BDT & b-tag


SUSY & DM Searches @ LHC O. Buchmüller
m_{LSP} [GeV]
Direct squark
$m_{SUSY} = m_{\tilde{q}}$
$\tilde{q} \to q\chi_1^0$
CMS arXiv:1502.04358
$\tilde{u}_L \to u\chi_1^0$
CMS arXiv:1502.04358
$\tilde{b} \to b\chi_1^0$
ATLAS arXiv:1308.2631
$\tilde{t} \to t\chi_1^0$
ATLAS arXiv:1407.0583
Direct stop in "gap"
$m_{SUSY} = m_{\tilde{t}}$
$\tilde{t} \to c\chi_1^0$
ATLAS arXiv:1407.0608
$\tilde{t} \to bff'\chi_1^0$
ATLAS arXiv:1407.0583
$\tilde{t} \to Wb\chi_1^0$
CMS arXiv:1308.1586
Gluino mediated
$m_{SUSY} = m_{\tilde{g}}$
$\tilde{g} \to q\bar{q}\chi_1^0$
ATLAS arXiv:1405.7875
$\tilde{g} \to b\bar{b}\chi_1^0$
CMS arXiv:1502.00300
$\tilde{g} \to t\bar{t}\chi_1^0$
ATLAS arXiv:1407.0600
$m_{SUSY} - m_{LSP} < m_t$
$m_{SUSY} - m_{LSP} < 2m_t$
$m_{\tilde{t}} - m_b - m_{LSP} < m_W$
$m_{\tilde{t}} - m_{LSP} < m_t$
BR=100%
all limits are observed nominal 95% CLs limits
RP conserved
m_{SUSY} [GeV]
0
250
500
750
1000
1250
1500


SUSY & DM Searches @ LHC O. Buchmüller
m_{LSP} [GeV]
1000
250
0
0
250
Direct squark
$m_{SUSY} = m_{\tilde{q}}$
$\tilde{q} \to q\chi_1^0$
CMS arXiv:1502.04358
$\tilde{u}_L \to u\chi_1^0$
CMS arXiv:1502.04358
$\tilde{b} \to b\chi_1^0$
ATLAS arXiv:1308.2631
Direct stop in "gap"
$m_{SUSY} = m_{\tilde{t}}$
$\tilde{t} \to c\chi_1^0$
ATLAS arXiv:1407.
$\tilde{t} \to bff'\chi_1^0$
ATLAS arXiv:1407.
Xiv:1308.1
Gluino mediated
$m_{SUSY} = m_{\tilde{g}}$
iv:1405.7875
:1502.00300
iv:1407.0600
ATLAS
$\int Ldt = 20.3\ fb^{-1}$, $\sqrt{s} = 8$ TeV
Observed limit ($\pm 1\ \sigma^{SUSY}_{theory}$)
Expected limit ($\pm 1\ \sigma_{exp}$)
LEP2 $\tilde{\mu}_R$ excluded
All limits at 95% CL
$m(\tilde{l}) < m(\tilde{\chi}_1^0)$
$m_{\tilde{\chi}_1^0}$ [GeV]
$m_{\tilde{l}_1}$ [GeV]
p
p
$\tilde{\ell}$
$\tilde{\ell}$
ℓ
ℓ
$\tilde{\chi}_1^0$
$\tilde{\chi}_1^0$
gluino/ squarks
charginos/ neutralinos
Higgs sector
sleptons
LSP
Mass / GeV
ATLAS arXiv:1403.5294
Signature
2 lepton + E_T^{miss}


SUSY & DM Searches @ LHC O. Buchmüller
m_{LSP} [GeV]
Direct squark
$m_{SUSY} = m_{\tilde{q}}$
$\tilde{q} \to q\chi_1^0$
CMS arXiv:1502.04358
$\tilde{u}_L \to u\chi_1^0$
CMS arXiv:1502.04358
$\tilde{b} \to b\chi_1^0$
ATLAS arXiv:1308.2631
$\tilde{t} \to t\chi_1^0$
ATLAS arXiv:1407.0583
Direct stop in "gap"
$m_{SUSY} = m_{\tilde{t}}$
$\tilde{t} \to c\chi_1^0$
ATLAS arXiv:1407.0608
$\tilde{t} \to bff'\chi_1^0$
ATLAS arXiv:1407.0583
$\tilde{t} \to Wb\chi_1^0$
CMS arXiv:1308.1586
Gluino mediated
$m_{SUSY} = m_{\tilde{g}}$
$\tilde{g} \to q\bar{q}\chi_1^0$
ATLAS arXiv:1405.7875
$\tilde{g} \to b\bar{b}\chi_1^0$
CMS arXiv:1502.00300
$\tilde{g} \to t\bar{t}\chi_1^0$
ATLAS arXiv:1407.0600
Direct slepton
$m_{SUSY} = m_{\tilde{l}}$
$\tilde{l}_R \to l^{\pm}\chi_1^0$
ATLAS: arXiv:1403.5294
$\tilde{l}_L \to l^{\pm}\chi_1^0$
ATLAS: arXiv:1403.5294
$m_{SUSY} - m_{LSP} < m_t$
$m_{SUSY} - m_{LSP} < 2m_t$
$m_{\tilde{t}} - m_b - m_{LSP} < m_W$
$m_{\tilde{t}} - m_{LSP} < m_t$
BR=100%
all limits are observed nominal 95% CLs limits
RP conserved
m_{SUSY} [GeV]
0
250
500
750
1000
1250
1500

Direct chargino/neutralino production

$X_2^0 X_1^{+-}$ production







heavy slepton"hard(er)"

Add $Z(l^+l^-)$+2jets topology in bins of E_t^{miss} to increase sensitivity for "heavy" slepton case



$X_1^+ X_1^-$ production

light slepton "easy"





ATLAS arXiv:1403.5294


SUSY & DM Searches @ LHC O. Buchmüller
m_{LSP} [GeV]
Direct squark
$m_{SUSY} = m_{\tilde{q}}$
$\tilde{q} \to q\chi_1^0$
CMS arXiv:1502.04358
$\tilde{u}_L \to u\chi_1^0$
CMS arXiv:1502.04358
$\tilde{b} \to b\chi_1^0$
ATLAS arXiv:1308.2631
$\tilde{t} \to t\chi_1^0$
ATLAS arXiv:1407.0583
Direct stop in "gap"
$m_{SUSY} = m_{\tilde{t}}$
$\tilde{t} \to c\chi_1^0$
ATLAS arXiv:1407.0608
$\tilde{t} \to bff'\chi_1^0$
ATLAS arXiv:1407.0583
$\tilde{t} \to Wb\chi_1^0$
CMS arXiv:1308.1586
Gluino mediated
$m_{SUSY} = m_{\tilde{g}}$
$\tilde{g} \to q\bar{q}\chi_1^0$
ATLAS arXiv:1405.7875
$\tilde{g} \to b\bar{b}\chi_1^0$
CMS arXiv:1502.00300
$\tilde{g} \to t\bar{t}\chi_1^0$
ATLAS arXiv:1407.0600
Direct slepton
$m_{SUSY} = m_{\tilde{l}}$
$\tilde{l}_R \to l^{\pm}\chi_1^0$
ATLAS: arXiv:1403.5294
$\tilde{l}_L \to l^{\pm}\chi_1^0$
ATLAS: arXiv:1403.5294
$m_{SUSY} - m_{LSP} < m_t$
$m_{SUSY} - m_{LSP} < 2m_t$
$m_{\tilde{t}} - m_b - m_{LSP} < m_W$
$m_{\tilde{t}} - m_{LSP} < m_t$
Direct $\chi_1^{\pm}/\chi_2^0$
$m_{SUSY} = m_{\chi_1^{\pm}} = m_{\chi_2^0}$
$\chi_1^{\pm}\chi_2^0$(light $\tilde{l}$)
CMS: arXiv:1405.7570
$\chi_1^{\pm}\chi_2^0$(heavy $\tilde{l}$)
CMS: arXiv:1405.7570
BR=100%
all limits are observed nominal 95% CLs limits
RP conserved
m_{SUSY} [GeV]
0
250
500
750
1000
1250
1500


SUSY & DM Searches @ LHC O. Buchmüller
m_{LSP} [GeV]
Direct squark
$m_{SUSY} = m_{\tilde{q}}$
$\tilde{q} \to q\chi_1^0$
CMS arXiv:1502.04358
$\tilde{u}_L \to u\chi_1^0$
CMS arXiv:1502.04358
$\tilde{b} \to b\chi_1^0$
ATLAS arXiv:1308.2631
$\tilde{t} \to t\chi_1^0$
ATLAS arXiv:1407.0583
Direct stop in "gap"
$m_{SUSY} = m_{\tilde{t}}$
$\tilde{t} \to c\chi_1^0$
ATLAS arXiv:1407.0608
$\tilde{t} \to bff'\chi_1^0$
ATLAS arXiv:1407.0583
$\tilde{t} \to Wb\chi_1^0$
CMS arXiv:1308.1586
Gluino mediated
$m_{SUSY} = m_{\tilde{g}}$
$\tilde{g} \to q\bar{q}\chi_1^0$
ATLAS arXiv:1405.7875
$\tilde{g} \to b\bar{b}\chi_1^0$
CMS arXiv:1502.00300
$\tilde{g} \to t\bar{t}\chi_1^0$
ATLAS arXiv:1407.0600
Direct slepton
$m_{SUSY} = m_{\tilde{l}}$
$\tilde{l}_R \to l^{\pm}\chi_1^0$
ATLAS: arXiv:1403.5294
$\tilde{l}_L \to l^{\pm}\chi_1^0$
ATLAS: arXiv:1403.5294
$m_{SUSY} - m_{LSP} < m_t$
$m_{SUSY} - m_{LSP} < 2m_t$
$m_{\tilde{t}} - m_b - m_{LSP} < m_W$
$m_{\tilde{t}} - m_{LSP} < m_t$
Direct $\chi_1^+\chi_1^-$
$m_{SUSY} = m_{\chi_1^{\pm}}$
$\chi_1^+\chi_1^-$(heavy $\tilde{l}$)
ATLAS arXiv:1403.5294
$\chi_1^+\chi_1^-$(light $\tilde{l}$)
ATLAS arXiv:1403.5294
BR=100%
all limits are observed nominal 95% CLs limits
RP conserved
Direct $\chi_1^{\pm}/\chi_2^0$
$m_{SUSY} = m_{\chi_1^{\pm}} = m_{\chi_2^0}$
$\chi_1^{\pm}\chi_2^0$(light $\tilde{l}$)
CMS: arXiv:1405.7570
$\chi_1^{\pm}\chi_2^0$(heavy $\tilde{l}$)
CMS: arXiv:1405.7570
m_{SUSY} [GeV]
0
250
500
750
1000
1250
1500

MASTERCODE

Resolving tension (g-2) and LHC



	X^2/n_{dof}	p-value
CMSSM	32.8/24	11 %
NUHM1	31.1/23	12 %

Can adding extra parameters resolve the tension between (g-2) and jets+MET constraints?

From MasterCode papers:
1312.5250, 1408.4060 and 1504.03260

Resolving tension (g-2) and LHC



	X^2/n_{dof}	p-value
CMSSM	32.8/24	11 %
NUHM1	31.1/23	12 %
NUHM2	30.3/22	11 %

NUHM2 can get (g-2) right but only at the expense of M_h and jets + MET constraints.

From MasterCode papers:
1312.5250, 1408.4060 and 1504.03260

Resolving tension (g-2) and LHC



	X^2/n_{dof}	p-value
CMSSM	32.8/24	11 %
NUHM1	31.1/23	12 %
NUHM2	30.3/22	11 %
pMSSM10	**20.5/18**	**31 %**

pMSSM10 resolves the tension between (g-2) and LHC constraints. This significantly improves the fit.

From MasterCode papers:
1312.5250, 1408.4060 and 1504.03260

CMSSM



From MasterCode papers:
1312.5250, 1408.4060 and 1504.03260

NUHM1



From MasterCode papers:
1312.5250, 1408.4060 and 1504.03260

NUHM2



From MasterCode papers: 1312.5250, 1408.4060 and 1504.03260

MasterCode: The two worlds of SUSY models

"Soft scale" ⟷ "GUT scale"

pMSSM10

M_1, M_2, M_3, $m_{\tilde{q}_{12}}$, $m_{\tilde{q}_3}$, $m_{\tilde{\ell}}$, A, M_A, $\tan\beta$, μ



CMSSM

$m_0,\ m_{1/2},$

$A_0,\ \tan\beta$

NUHM1

$m^2_{H_u} = m^2_{H_d}$

NUHM2

$m^2_{H_u} \neq m^2_{H_d}$

CMSSM Today: Mq-Mg Search plane



From MasterCode papers:
1312.5250, 1408.4060 and 1504.03260

CMSSM Today: Mq-Mg Search plane



From MasterCode papers:
1312.5250, 1408.4060 and 1504.03260

CMSSM Today: Mq-Mg Search plane



From MasterCode papers:
1312.5250, 1408.4060 and 1504.03260

CMSSM Today: Mq-Mg Search plane



Models in Comparison in “Mq-Mg Search plane”



From MasterCode papers:
1312.5250, 1408.4060 and 1504.03260

pMSSM10: parameter space



SUSY PROJECTION OF DIFFICULT CHANNELS



SUSY & DM Searches @ LHC O. Buchmüller


SUSY & DM Searches @ LHC O. Buchmüller
m_{LSP} [GeV]
1000
500
250
0
LHC: 8 TeV 20 fb^{-1}
LHC: 14 TeV 300 fb^{-1}
Direct squark
$m_{SUSY} = m_{\tilde{q}}$
$\tilde{t} \to t\chi_1^0$ ATLAS-CONF-2013-037
Direct slepton
$\tilde{l}_R \to l^{\pm}\chi_1^0$ ATLAS-CONF-2013-049
Direct $\chi_1^{\pm}/\chi_2^0$
$\chi_1^{\pm}\chi_2^0(heavy\ \tilde{l})$
CMS-PAS-SUS-13-006
$m_{SUSY} = m_{\chi_1^{\pm}} = m_{\chi_2^0}$
BR=100%
all limits are observed nominal 95% CLs limits RP conserved
m_{SUSY} [GeV]
0
250
500
750
1000
1250
1500


SUSY & DM Searches @ LHC O. Buchmüller
m_{LSP} [GeV]
1000
500
250
0
0
250
500
750
1000
1250
1500
m_{SUSY} [GeV]
LHC: 8 TeV 20 fb^{-1}
LHC: 14 TeV 300 fb^{-1}
HL-LHC: 14 TeV 3000 fb^{-1}
Direct squark
$m_{SUSY} = m_{\tilde{q}}$
$\tilde{t} \to t\chi_1^0$ ATLAS-CONF-2013-037
Direct slepton
$\tilde{l}_R \to l^{\pm}\chi_1^0$ ATLAS-CONF-2013-049
Direct $\chi_1^{\pm}/\chi_2^0$
$\chi_1^{\pm}\chi_2^0(heavy\ \tilde{l})$
CMS-PAS-SUS-13-006
$m_{SUSY} = m_{\chi_1^{\pm}} = m_{\chi_2^0}$
BR=100%
all limits are observed nominal 95% CLs limits
RP conserved


SUSY & DM Searches @ LHC O. Buchmüller
m_{LSP} [GeV]
1000
500
250
0
0
250
500
750
1000
1250
1500
m_{SUSY} [GeV]
LHC: 8 TeV 20 fb^{-1}
LHC: 14 TeV 300 fb^{-1}
HL-LHC: 14 TeV 3000 fb^{-1}
Direct squark
$m_{SUSY} = m_{\tilde{q}}$
$\tilde{t} \to t\chi_1^0$ ATLAS-CONF-2013-037
Direct slepton
$\tilde{l}_R \to l^{\pm}\chi_1^0$ ATLAS-CONF-2013-049
Direct $\chi_1^{\pm}/\chi_2^0$
$\chi_1^{\pm}\chi_2^0(heavy\ \tilde{l})$
CMS-PAS-SUS-13-006
$m_{SUSY} = m_{\chi_1^{\pm}} = m_{\chi_2^0}$
BR=100%
all limits are observed nominal 95% CLs limits RP conserved


SUSY & DM Searches @ LHC O. Buchmüller
m_{LSP} [GeV]
1000
500
250
0
0
250
500
750
1000
1250
1500
m_{SUSY} [GeV]
LHC: 8 TeV 20 fb^{-1}
LHC: 14 TeV 300 fb^{-1}
HL-LHC: 14 TeV 3000 fb^{-1}
Direct squark
$m_{SUSY} = m_{\tilde{q}}$
$\tilde{t} \to t\chi_1^0$ ATLAS-CONF-2013-037
Direct slepton
$\tilde{l}_R \to l^{\pm}\chi_1^0$ ATLAS-CONF-2013-049
Direct $\chi_1^{\pm}/\chi_2^0$
$\chi_1^{\pm}\chi_2^0(heavy\ \tilde{l})$
CMS-PAS-SUS-13-006
$m_{SUSY} = m_{\chi_1^{\pm}} = m_{\chi_2^0}$
1 TeV
linear
collider
BR=100%
all limits are
observed nominal
95% CLs limits
RP conserved

EFT VALIDITY REGION ILLUSTRATED

Effective Field Theory (EFT) Interpretation

Example of considered operators:



$$O_V = \frac{(\bar{\chi}\gamma_\mu\chi)(\bar{q}\gamma_\mu q)}{\Lambda^2}$$ Vector operator, s-channel

$$O_{AV} = \frac{(\bar{\chi}\gamma_\mu\gamma_5\chi)(\bar{q}\gamma_\mu\gamma_5 q)}{\Lambda^2}$$ Axial vector operator, s-channel

Assumption of EFT

If the operator (e.g. V or AV) mediator is suitably(!!) heavy it can be integrated out to obtain the effective V or AV contact operator. In this case (and only this case), the contact interaction scale Λ is related to the parameters entering the Lagrangian:

$$\Lambda = \frac{M_{mediator}}{\sqrt{g_q g_\chi}}$$ (relation in the full theory)

Validity of Effective Field Theory Limits

Recent work from OB, M.Dolan,C.McCabe: arXiv:1308.6799

- Compare Effective Field Theory (EFT) with Full Theory (FT)

EFT approach





FT
one diagram
"simplified model"

Use vector and axial-vector mediators (e.g. Z') as example - scalar are similar in conclusion!



Compare prediction of FT with EFT in m_{med} – m_{DM} plane.
Three regions become visible:

Region I: EFT and FT agree better then 20%

- EFT is valid!

Region II: EFT yields significant weaker limits then FT

- EFT limits are too conservative!

Region III: EFT yields significant stronger limits then FT

- EFT limits are too aggressive!

Validity of Effective Field Theory Limits

Recent work from OB, M.Dolan,C.McCabe: arXiv:1308.6799

- Compare Effective Field Theory (EFT) with Full Theory (FT)

EFT approach





FT
one diagram
"simplified model"

Use vector and axial-vector mediators (e.g. Z') as example - scalar are similar in conclusion!



Three Regions as function of mediator mass:

Region I: Heavy m_{med}

- EFT is valid!

Region II: Medium m_{med} – Resonant enhancement

- EFT limits are too conservative!

Region III: Low m_{med}

- EFT limits are too aggressive!

Validity of Effective Field Theory Limits

Recent work from OB, M.Dolan,C.McCabe: arXiv:1308.6799

- Compare Effective Field Theory (EFT) with Full Theory (FT)

EFT approach





FT
one diagram
"simplified model"

Use v … clusion!

Conclusion:
The EFT is not an appropriate framework for a comprehensive Interpretation of DM searches at colliders and especially must taken with very (as in VERY) special care when comparing with other experiments such as Direct Detection!



Region I: Heavy m_{med}
- EFT is valid!

Region II: Medium m_{med} – Resonant enhancement
- EFT limits are too conservative!

Region III: Low m_{med}
- EFT limits are too aggressive!

What those this imply on model-dependences of EFT limits?



Look at EFT validity in m_{DM} – coupling* plane!

* Coupling chose such that CMS EFT limit on Λ applies to FT

Model-dependences of EFT limits



Look at EFT validity in m_{DM} – coupling* plane!

1. Region in which EFT is valid

For this we calculate the minimum coupling

$$\sqrt{g_q g_\chi} = m_{med}/\Lambda_{CMS}$$

that the simplified model must have for the EFT limits to apply. This is defined by region I (i.e. better then 20% agreement of FT and EFT).

* Coupling chose such that CMS EFT limit on Λ applies to FT

Model-dependences of EFT limits



Look at EFT validity in m_{DM} – coupling* plane!

1. Region in which EFT is valid (20%)
2. Require compatibility with relic density

When exclude the region in which relic abundance is larger then the observed value of $\Omega_{\chi\chi}h^2 = 0.119$ only mediator masses above a few hundred GeV fulfill this.

* Coupling chose such that CMS EFT limit on Λ applies to FT

Model-dependences of EFT limits



Look at EFT validity in m_{DM} – coupling* plane!

1. Must require $m_{med} < \Gamma_{med}$
2. Region in which EFT is valid (20%)
3. Require compatibility with relic density
4. Require theory to be perturbative (<4π)

When we also require that the region/theory must be perturbative:

$$\sqrt{g_q g_\chi} < 4\pi$$

only a very small region is left!

EFT limits of monojet searches only apply to a very (as in VERY) small class of DM models!

Model-dependences of EFT limits



Look at EFT validity in m_{DM} – coupling* plane!

1. Region in which EFT is valid (20%)
2. Require compatibility with relic density
3. Require theory to be perturbative (<4π)
4. $m_{med} < \Gamma_{med}$ ALWAYS!

We also find that for all DM models the EFT Is valid the mass of the mediator must be Smaller than its width!

In the reaming part of the plot:

$$\sqrt{g_q g_\chi} > 2$$

a particle-like interpretation of the mediator is doubtful because of $m_{med} < \Gamma_{med}$!

See discussion about equation 3.5 in arXiv:1308.6799 for further details.

What those this imply on model-dependences of EFT limits?



Look at EFT validity in m_{DM} – coupling* plane!

1. Region in which EFT is valid (20%)
2. Require compatibility with relic density
3. Require theory to be perturbative (<4π)
4. $m_{med} < \Gamma_{med}$ ALWAYS!

The observation that all DM theories for which the EFT is valid must have $m_{med} < \Gamma_{med}$ and the small class to models it applies in any case leads to the conclusion the EFT only applies to a very small class of DM models.
EFT limits of monojet searches are therefore highly model-depended!

Collider vs Direct Detection

M_{DM}	M_{med}
g_q	g_{DM}

Vector



Axial vector



arXiV:1407.8257

Mono-Mania (at the LHC)

Mono-Z



Mono-photon



Mono-jet



Mono-Higgs



Mono-W



Mono-top



Mono-Mania (at the LHC)

Mono-Z



Mono-photon



Mono-jet



Mono-Higgs



Mono-W



Mono-top