New Directions in Searching for the Dark Universe

Surjeet Rajendran,
UC Berkeley

Dark Matter

Dark Matter



A New Particle

Non gravitational interactions?

Dark Matter



A New Particle

Non gravitational interactions?

How do we detect them?

Weak effects. Need high precision

Precision Instruments

Impressive developments in the past two decades



$$\text{Magnetic Field} \lesssim 10^{-16} \frac{\text{T}}{\sqrt{\text{Hz}}}$$

(SQUIDs, atomic magnetometers)



$$\text{Accelerometers} \lesssim 10^{-13} \frac{\text{g}}{\sqrt{\text{Hz}}}$$

(atom and optical interferometers)

Precision Instruments

Impressive developments in the past two decades





$$\text{Magnetic Field} \lesssim 10^{-16} \frac{\text{T}}{\sqrt{\text{Hz}}}$$

(SQUIDs, atomic magnetometers)

$$\text{Accelerometers} \lesssim 10^{-13} \frac{\text{g}}{\sqrt{\text{Hz}}}$$

(atom and optical interferometers)

Rapid technological advancements

Use to detect new physics?

Outline

1. Brief Theory Overview

2. Axion Detection with Nuclear Magnetic Resonance

3. Dark Photon Detection with Radios

4. Dark Matter Detection with Accelerometers

5. Conclusions

The Dark Matter Landscape

The Dark Matter Landscape



Fit in galaxy

The Dark Matter Landscape



Fit in galaxy

Standard Model scale ~ 100 GeV

The Dark Matter Landscape



Fit in galaxy

Standard Model scale ~ 100 GeV

One Possibility: Same scale for Dark Matter?
Weakly Interacting Massive Particles (WIMPs)

The Dark Matter Landscape



Fit in galaxy

Standard Model scale ~ 100 GeV

One Possibility: Same scale for Dark Matter?
Weakly Interacting Massive Particles (WIMPs)

Other Generic Candidates: Axions, Massive Vector Bosons

The Dark Matter Landscape



Fit in galaxy

Standard Model scale ~ 100 GeV

One Possibility: Same scale for Dark Matter?
Weakly Interacting Massive Particles (WIMPs)

Other Generic Candidates: Axions, Massive Vector Bosons

How do we search for them?
This Talk: Bosons between GHz - Hz
Range includes popular candidates such as the QCD axion

Bosonic Dark Matter

Photons

$$\vec{E} = E_0 \cos(\omega t - \omega x)$$

Detect Photon by measuring time varying field

Bosonic Dark Matter

Photons

$$\vec{E} = E_0 \cos(\omega t - \omega x)$$

Detect Photon by measuring time varying field

Dark Bosons

Early Universe:
Misalignment Mechanism



$$a(t) \sim a_0 \cos(m_a t)$$

Spatially uniform, oscillating field

$$m_a^2 a_0^2 \sim \rho_{DM}$$

Bosonic Dark Matter

Photons

$$\vec{E} = E_0 \cos(\omega t - \omega x)$$

Detect Photon by measuring time varying field

Dark Bosons

Early Universe: Misalignment Mechanism



$$a(t) \sim a_0 \cos(m_a t)$$

Spatially uniform, oscillating field

$$m_a^2 a_0^2 \sim \rho_{DM}$$

Today: Random Field



Correlation length
~ $1/(m_a v)$

Coherence Time
~ $1/(m_a v^2)$
~ 1 s (MHz/m_a)

Bosonic Dark Matter

Photons

$$\vec{E} = E_0 \cos(\omega t - \omega x)$$

Detect Photon by measuring time varying field

Dark Bosons

Early Universe: Misalignment Mechanism



$$a(t) \sim a_0 \cos(m_a t)$$

Spatially uniform, oscillating field

$$m_a^2 a_0^2 \sim \rho_{DM}$$

Today: Random Field



Correlation length
~ $1/(m_a v)$

Coherence Time
~ $1/(m_a v^2)$
~ 1 s (MHz/m_a)

Detect effects of oscillating dark matter field

Resonance possible. Q ~ 10^6 (set by v ~ 10^{-3})

What kind of Bosons?

Naturalness. Structure set by symmetries.

What kind of Bosons?

Naturalness. Structure set by symmetries.

Spin 0

Axions or ultra weak coupling

Many UV theories

What kind of Bosons?

Naturalness. Structure set by symmetries.

Spin 0

Axions or ultra weak coupling

Many UV theories

E&M	QCD	Spin	Higgs
$\left(\frac{a}{f_a} F\tilde{F}\right)$	$\left(\frac{a}{f_a} G\tilde{G}\right)$	$\left(\frac{\partial_\mu a}{f_a} \bar{N}\gamma^\mu\gamma_5 N\right)$	$\left(g\phi H^2\right)$
	QCD Axion	General Axions	Higgs Portal/ Relaxion

What kind of Bosons?

Naturalness. Structure set by symmetries.

Spin 0

Axions or ultra weak coupling

Many UV theories

Spin 1

Anomaly free

Standard Model couplings

E&M	QCD	Spin	Higgs
$\left(\frac{a}{f_a} F\tilde{F}\right)$	$\left(\frac{a}{f_a} G\tilde{G}\right)$	$\left(\frac{\partial_\mu a}{f_a} \bar{N}\gamma^\mu\gamma_5 N\right)$	$\left(g\phi H^2\right)$
	QCD Axion	General Axions	Higgs Portal/ Relaxion

What kind of Bosons?

Naturalness. Structure set by symmetries.

Spin 0

Axions or ultra weak coupling

Many UV theories

Spin 1

Anomaly free

Standard Model couplings

E&M	QCD	Spin	Higgs	Spin	E&M	Current
$\left(\frac{a}{f_a} F\tilde{F}\right)$	$\left(\frac{a}{f_a} G\tilde{G}\right)$	$\left(\frac{\partial_\mu a}{f_a} \bar{N}\gamma^\mu\gamma_5 N\right)$	$\left(g\phi H^2\right)$	$\left(\frac{F'_{\mu\nu}}{f_a} \bar{N}\sigma^{\mu\nu} N\right)$	$\left(\epsilon F' F\right)$	$\left(g A'_\mu J^\mu_{B-L}\right)$
	QCD Axion	General Axions	Higgs Portal/ Relaxion	Dipole moment	Kinetic Mixing	B-L

What kind of Bosons?

Naturalness. Structure set by symmetries.

Spin 0

Axions or ultra weak coupling

Many UV theories

Spin 1

Anomaly free

Standard Model couplings

E&M	QCD	Spin	Higgs	Spin	E&M	Current
$\left(\frac{a}{f_a} F\tilde{F}\right)$	$\left(\frac{a}{f_a} G\tilde{G}\right)$	$\left(\frac{\partial_\mu a}{f_a} \bar{N}\gamma^\mu\gamma_5 N\right)$	$\left(g\phi H^2\right)$	$\left(\frac{F'_{\mu\nu}}{f_a} \bar{N}\sigma^{\mu\nu} N\right)$	$\left(\epsilon F' F\right)$	$\left(g A'_\mu J^\mu_{B-L}\right)$
	QCD Axion	General Axions	Higgs Portal/ Relaxion	Dipole moment	Kinetic Mixing	B-L

$$\text{Dark Matter} \implies a = a_0 \cos\left(m_a t\right)$$

$$\text{Hz} \lessapprox m_a \lessapprox \text{GHz}$$

What kind of Bosons?

Naturalness. Structure set by symmetries.

Spin 0

Axions or ultra weak coupling

Many UV theories

Spin 1

Anomaly free

Standard Model couplings

E&M	QCD	Spin	Higgs	Spin	E&M	Current
$\left(\frac{a}{f_a} F\tilde{F}\right)$	$\left(\frac{a}{f_a} G\tilde{G}\right)$	$\left(\frac{\partial_\mu a}{f_a} \bar{N}\gamma^\mu\gamma_5 N\right)$	$\left(g\phi H^2\right)$	$\left(\frac{F'_{\mu\nu}}{f_a} \bar{N}\sigma^{\mu\nu} N\right)$	$\left(\epsilon F' F\right)$	$\left(g A'_\mu J^\mu_{B-L}\right)$
Current Searches (m_a ~ GHz)	QCD Axion	General Axions	Higgs Portal/ Relaxion	Dipole moment	Kinetic Mixing	B-L

$$\text{Dark Matter} \implies a = a_0 \cos\left(m_a t\right)$$

$$\text{Hz} \lessapprox m_a \lessapprox \text{GHz}$$

What kind of Bosons?

Naturalness. Structure set by symmetries.

Spin 0

Axions or ultra weak coupling

Many UV theories

Spin 1

Anomaly free

Standard Model couplings

E&M	QCD	Spin	Higgs	Spin	E&M	Current
$\left(\frac{a}{f_a} F\tilde{F}\right)$	$\left(\frac{a}{f_a} G\tilde{G}\right)$	$\left(\frac{\partial_\mu a}{f_a} \bar{N}\gamma^\mu\gamma_5 N\right)$	$\left(g\phi H^2\right)$	$\left(\frac{F'_{\mu\nu}}{f_a} \bar{N}\sigma^{\mu\nu} N\right)$	$\left(\epsilon F' F\right)$	$\left(g A'_\mu J^\mu_{B-L}\right)$
Current Searches (m_a ~ GHz)	QCD Axion	General Axions	Higgs Portal/ Relaxion	Dipole moment	Kinetic Mixing	B-L

$$\text{Dark Matter} \implies a = a_0 \cos\left(m_a t\right)$$

This Talk

$$\boxed{\text{Hz} \lessapprox m_a \lessapprox \text{GHz}}$$

SIMONS FOUNDATION

HEISING - SIMONS FOUNDATION

DFG Deutsche Forschungsgemeinschaft

Cosmic Axion Spin Precession Experiment (CASPEr)

with

Dmitry Budker
Peter Graham
Micah Ledbetter
Alex Sushkov

PRX **4** (2014) arXiv: 1306.6089
PRD **88** (2013) arXiv: 1306.6088
PRD **84** (2011) arXiv: 1101.2691

CASPEr: Axion Effects on Spin

CASPEr: Axion Effects on Spin

General Axions

Neutron



CASPEr: Axion Effects on Spin

General Axions

Neutron in
Axion Wind



$$H_N \supset \frac{a}{f_a}\vec{v_a}.\vec{S}_N$$

Spin rotates about
dark matter velocity

CASPEr: Axion Effects on Spin

General Axions

Neutron in
Axion Wind



$$H_N \supset \frac{a}{f_a}\vec{v_a}.\vec{S}_N$$

Spin rotates about
dark matter velocity

CASPEr: Axion Effects on Spin

General Axions

Neutron in
Axion Wind



$$H_N \supset \frac{a}{f_a}\vec{v_a}.\vec{S}_N$$

Spin rotates about
dark matter velocity

Effective time varying
magnetic field

$$B_{eff} \lessapprox 10^{-16}\ \cos\left(m_a t\right)\ \mathrm{T}$$

CASPEr: Axion Effects on Spin

General Axions

Neutron in
Axion Wind



$$H_N \supset \frac{a}{f_a}\vec{v_a}.\vec{S}_N$$

Spin rotates about
dark matter velocity

Effective time varying
magnetic field

$$B_{eff} \lessapprox 10^{-16}\ \cos\left(m_a t\right)\ \mathrm{T}$$

Other light dark matter (e.g. dark photons) also
induce similar spin precession

CASPEr: Axion Effects on Spin

General Axions

Neutron in
Axion Wind

$\left(\frac{\partial_\mu a}{f_a}\bar{N}\gamma^\mu\gamma_5 N\right)$



$$H_N \supset \frac{a}{f_a}\vec{v_a}.\vec{S}_N$$

Spin rotates about
dark matter velocity

Effective time varying
magnetic field

$$B_{eff} \lessapprox 10^{-16}\ \cos\left(m_a t\right)\ \mathrm{T}$$

Other light dark matter (e.g. dark photons) also
induce similar spin precession

QCD Axion

Neutron



CASPEr: Axion Effects on Spin

General Axions

Neutron in
Axion Wind

$\left(\frac{\partial_\mu a}{f_a}\bar{N}\gamma^\mu\gamma_5 N\right)$



$$H_N \supset \frac{a}{f_a}\vec{v_a}.\vec{S}_N$$

Spin rotates about
dark matter velocity

Effective time varying
magnetic field

$$B_{eff} \lessapprox 10^{-16}\,\cos\left(m_a t\right)\,\mathrm{T}$$

Other light dark matter (e.g. dark photons) also
induce similar spin precession

QCD Axion

Neutron in
QCD Axion Dark Matter



$\left(\frac{a}{f_a}G\tilde{G}\right)$

QCD axion induces electric dipole moment
for neutron and proton

Dipole moment
along nuclear spin

Oscillating dipole: $d \sim 3\times 10^{-34}\,\cos\left(m_a t\right)\,e\,\mathrm{cm}$

CASPEr: Axion Effects on Spin

General Axions

Neutron in
Axion Wind



$$H_N \supset \frac{a}{f_a}\vec{v_a}.\vec{S}_N$$

Spin rotates about
dark matter velocity

Effective time varying
magnetic field

$$B_{eff} \lessapprox 10^{-16}\,\cos\left(m_a t\right)\,\mathrm{T}$$

Other light dark matter (e.g. dark photons) also
induce similar spin precession

QCD Axion

Neutron in
QCD Axion Dark Matter



QCD axion induces electric dipole moment
for neutron and proton

Dipole moment
along nuclear spin

Oscillating dipole: $d \sim 3\times 10^{-34}\,\cos\left(m_a t\right)\,e\,\mathrm{cm}$

Apply electric field, spin rotates

CASPEr: Axion Effects on Spin

General Axions

Neutron in
Axion Wind



$$H_N \supset \frac{a}{f_a}\vec{v_a}.\vec{S}_N$$

Spin rotates about
dark matter velocity

Effective time varying
magnetic field

$$B_{eff} \lessapprox 10^{-16}\,\cos\left(m_a t\right)\,\mathrm{T}$$

Other light dark matter (e.g. dark photons) also
induce similar spin precession

QCD Axion

Neutron in
QCD Axion Dark Matter



QCD axion induces electric dipole moment
for neutron and proton

Dipole moment
along nuclear spin

Oscillating dipole: $d \sim 3\times 10^{-34}\,\cos\left(m_a t\right)\,e\,\mathrm{cm}$

Apply electric field, spin rotates

CASPEr: Axion Effects on Spin

General Axions

Neutron in
Axion Wind



$$H_N \supset \frac{a}{f_a}\vec{v_a}.\vec{S}_N$$

Spin rotates about
dark matter velocity

Effective time varying
magnetic field

$$B_{eff} \lessapprox 10^{-16}\ \cos\left(m_a t\right)\ \mathrm{T}$$

Other light dark matter (e.g. dark photons) also
induce similar spin precession

Measure Spin
Rotation,
detect Axion

QCD Axion

Neutron in
QCD Axion Dark Matter



QCD axion induces electric dipole moment
for neutron and proton

Dipole moment
along nuclear spin

Oscillating dipole: $d \sim 3 \times 10^{-34}\ \cos\left(m_a t\right)\ e\,\mathrm{cm}$

Apply electric field, spin rotates

CASPEr

Axion affects physics of nucleus, NMR is sensitive probe



Larmor frequency = axion mass ➔ resonant enhancement

CASPEr

Axion affects physics of nucleus, NMR is sensitive probe



Larmor frequency = axion mass ➔ resonant enhancement

SQUID measures resulting transverse magnetization

NMR well established technology, noise understood, similar setup to previous experiments

Example materials: LXe, ferroelectric $PbTiO_3$, many others

CASPEr-General Axions



~ year to scan one decade of frequency

CASPEr-General Axions



~ year to scan one decade of frequency

CASPEr-QCD Axion



Verify signal with spatial coherence of axion field

Dark Photon Detection with a Radio

with

Peter Graham
Kent Irwin
Saptarshi Chaudhuri
Jeremy Mardon
Yue Zhao

arXiv: 1411.7382

Dark Photon Dark Matter

Many theories/vacua have additional, decoupled sectors, new U(1)'s

Natural coupling (dim. 4 operator): $\mathcal{L} \supset \varepsilon F F'$

mass basis:

$$\mathcal{L} = -\frac{1}{4}\left(F_{\mu\nu}F^{\mu\nu} + F'_{\mu\nu}F'^{\mu\nu}\right) + \frac{1}{2}m_{\gamma'}^2 A'_\mu A'^\mu - eJ^\mu_{EM}\left(A_\mu + \varepsilon A'_\mu\right)$$

photon with small mass and suppressed couplings to all charged particles

Dark Photon Dark Matter

Many theories/vacua have additional, decoupled sectors, new U(1)'s

Natural coupling (dim. 4 operator): $\mathcal{L} \supset \varepsilon F F'$

mass basis:

$$\mathcal{L} = -\frac{1}{4}\left(F_{\mu\nu}F^{\mu\nu} + F'_{\mu\nu}F'^{\mu\nu}\right) + \frac{1}{2}m_{\gamma'}^2 A'_\mu A'^\mu - eJ^\mu_{EM}\left(A_\mu + \varepsilon A'_\mu\right)$$

photon with small mass and suppressed couplings to all charged particles

oscillating E' field
(dark matter)

Dark Photon Dark Matter

Many theories/vacua have additional, decoupled sectors, new U(1)'s

Natural coupling (dim. 4 operator): $\mathcal{L} \supset \varepsilon F F'$

mass basis:

$$\mathcal{L} = -\frac{1}{4}\left(F_{\mu\nu}F^{\mu\nu} + F'_{\mu\nu}F'^{\mu\nu}\right) + \frac{1}{2}m_{\gamma'}^2 A'_\mu A'^\mu - eJ^\mu_{EM}\left(A_\mu + \varepsilon A'_\mu\right)$$

photon with small mass and suppressed couplings to all charged particles

oscillating E' field (dark matter)

can drive current behind EM shield

Dark Matter Radio Station



EXPECTED REACH

Stage 1: size ~50 cm, T= 4K, Q=10^6, 1 year scan



Stage 2: size ~1 m, T= 10mK, Q=10^6, 1 year scan

Dark Matter Detection with Accelerometers

(under development)

with

Peter Graham
David Kaplan
Jeremy Mardon
William Terrano

B-L Dark Matter

Other than electromagnetism, only other anomaly free standard model current

$$\mathcal{L} = -\frac{1}{4}\left(F'_{\mu\nu}F'^{\mu\nu}\right) + \frac{1}{2}m^2_{\gamma'}A'_\mu A'^\mu - gJ^\mu_{B-L}A'_\mu$$

Protons, Neutrons, Electrons and Neutrinos are all charged

Electrically neutral atoms are charged under B-L

Force experiments constrain $g < 10^{-21}$

B-L Dark Matter

Other than electromagnetism, only other anomaly free standard model current

$$\mathcal{L} = -\frac{1}{4}\left(F'_{\mu\nu}F'^{\mu\nu}\right) + \frac{1}{2}m^2_{\gamma'}A'_\mu A'^\mu - gJ^\mu_{B-L}A'_\mu$$

Protons, Neutrons, Electrons and Neutrinos are all charged

Electrically neutral atoms are charged under B-L

Force experiments constrain $g < 10^{-21}$

oscillating E' field (dark matter)

can accelerate atoms

Force depends on net neutron number - violates equivalence principle. Dark matter exerts time dependent equivalence principle violating force!

B-L Dark Matter

Acceleration Per Baryon: $\frac{gE'}{m_n} \sim 10^{-10}\,\frac{\mathrm{m}}{\mathrm{s}^2}\left(\frac{g}{10^{-21}}\right)$

Atomic Accelerometers $\gtrapprox 10^{-12}\,\frac{\mathrm{m}}{\mathrm{s}^2\sqrt{\mathrm{Hz}}}$ (@ 1 Hz)

Dark Matter force depends upon net neutron number

Time dependent equivalence principle violation!

B-L Dark Matter

Acceleration Per Baryon: $\frac{gE'}{m_n} \sim 10^{-10}\,\frac{\mathrm{m}}{\mathrm{s}^2}\left(\frac{g}{10^{-21}}\right)$

Atomic Accelerometers $\gtrsim 10^{-12}\,\frac{\mathrm{m}}{\mathrm{s}^2\sqrt{\mathrm{Hz}}}$ (@ 1 Hz)

Dark Matter force depends upon net neutron number
Time dependent equivalence principle violation!

Improvements possible with resonant schemes

Seems promising!

Stanford Test Facility

The Relaxion

$$\mathcal{L} \supset (-M^2 + g\phi)|h|^2 + gM^2\phi + g^2\phi^2 + \cdots + \Lambda^4 \cos\frac{\phi}{f}$$

Hierarchy problem solved through cosmic evolution - does not require any new physics at the LHC

ϕ is a light scalar coupled to higgs with small coupling g

$$\Longrightarrow \frac{g\phi}{v} m_q \bar{q} q$$

$$\text{Dark matter } \phi \implies \phi = \phi_0 \cos\left(m_\phi \left(t - \vec{\mathrm{v}}.\vec{x}\right)\right)$$

Time variation of masses of fundamental particles

$$\Longrightarrow \quad \text{force on atoms } \frac{g\nabla\phi}{v} m_q \sim \frac{g m_\phi \vec{\mathrm{v}}}{v} m_q$$

Force violates equivalence principle. Time dependent equivalence principle violation!

Conclusions

WIMPs

Scalable

Goodman & Witten (1985): $\sigma \sim 10^{-38}\,\mathrm{cm}^2$

Z

h

W



WIMPs

Scalable

Goodman & Witten (1985): $\sigma \sim 10^{-38}\,\mathrm{cm}^2$

Z

h

W



Similar approach seems possible in searching for oscillating fields

The Dark Matter Landscape



The Dark Matter Landscape



Search for single, hard particle scattering



The Dark Matter Landscape



Time dependent moments of coherent classical field

Interactions restricted by symmetry

Frequencies can naturally be lab accessible (Hz - GHz)

Lab-scale experiments

Search for single, hard particle scattering



The Dark Matter Landscape



Time dependent moments of coherent classical field

Interactions restricted by symmetry

Frequencies can naturally be lab accessible (Hz - GHz)

Lab-scale experiments

Search for single, hard particle scattering



How do we cover full range?

Backup

A Different Operator For Axion Detection

So how can we detect high f_a axions?

Strong CP problem: $\mathcal{L} \supset \theta\, G\widetilde{G}$ creates a nucleon EDM $d \sim 3 \times 10^{-16}\, \theta\, e\,\mathrm{cm}$

the axion: $\mathcal{L} \supset \dfrac{a}{f_a} G\widetilde{G}$ creates a nucleon EDM $d \sim 3 \times 10^{-16}\, \dfrac{a}{f_a}\, e\,\mathrm{cm}$

A Different Operator For Axion Detection

So how can we detect high f_a axions?

Strong CP problem: $\mathcal{L} \supset \theta\, G\widetilde{G}$ creates a nucleon EDM $d \sim 3 \times 10^{-16}\, \theta\, e\,\mathrm{cm}$

the axion: $\mathcal{L} \supset \dfrac{a}{f_a} G\widetilde{G}$ creates a nucleon EDM $d \sim 3 \times 10^{-16}\, \dfrac{a}{f_a}\, e\,\mathrm{cm}$

$$a(t) \sim a_0 \cos\left(m_a t\right) \quad \text{with} \quad m_a \sim \frac{(200\ \mathrm{MeV})^2}{f_a} \sim \mathrm{MHz}\left(\frac{10^{16}\ \mathrm{GeV}}{f_a}\right)$$

axion dark matter $\rho_{\mathrm{DM}} \sim m_a^2 a^2 \sim (200\mathrm{MeV})^4 \left(\dfrac{a}{f_a}\right)^2 \sim 0.3\, \dfrac{\mathrm{GeV}}{\mathrm{cm}^3}$

so today: $\left(\dfrac{a}{f_a}\right) \sim 3 \times 10^{-19}$ independent of f_a

axion gives all nucleons an oscillating EDM (kHz-GHz) independent of f_a,
a non-derivative operator

Constraints and Searches



Constraints and Searches



Constraints and Searches



in most models: $\mathcal{L} \supset \frac{a}{f_a} F\widetilde{F} = \frac{a}{f_a} \vec{E} \cdot \vec{B}$

laser experiments:



axion emission affects SN1987A, White Dwarfs, other astrophysical objects

collider & laser experiments, ALPS, CAST

Constraints and Searches



Axion dark matter



in most models: $\mathcal{L} \supset \frac{a}{f_a} F\widetilde{F} = \frac{a}{f_a}\vec{E}\cdot\vec{B}$

axion-photon conversion suppressed $\propto \frac{1}{f_a^2}$

size of cavity increases with f_a

signal $\propto \frac{1}{f_a^3}$

microwave cavity (ADMX)

laser experiments:



$\propto \frac{1}{f_a^4}$

axion emission affects SN1987A, White Dwarfs, other astrophysical objects

collider & laser experiments, ALPS, CAST

Constraints and Searches



in most models: $\mathcal{L} \supset \frac{a}{f_a} F\widetilde{F} = \frac{a}{f_a}\vec{E}\cdot\vec{B}$

axion-photon conversion suppressed $\propto \frac{1}{f_a^2}$

size of cavity increases with f_a

signal $\propto \frac{1}{f_a^3}$

laser experiments:



$\propto \frac{1}{f_a^4}$

axion emission affects SN1987A, White Dwarfs, other astrophysical objects

collider & laser experiments, ALPS, CAST

Other ways to search for light (high f_a) axions?

Axions and the CMB

Assuming BICEP detected gravitational waves in the CMB (some tension with Planck):

$$H_{\rm inf} \sim 10^{14}\ {\rm GeV}$$

if symmetry broken after inflation → topological defects (strings + domain walls), constrained by observations

if symmetry broken before inflation → inflation can induce isocurvature perturbations of axion, weak constraint on ALPs probed by CASPEr.

Axions and the CMB

Assuming BICEP detected gravitational waves in the CMB (some tension with Planck):

$$H_{\rm inf} \sim 10^{14}\ {\rm GeV}$$

if symmetry broken after inflation → topological defects (strings + domain walls), constrained by observations

if symmetry broken before inflation → inflation can induce isocurvature perturbations of axion, weak constraint on ALPs probed by CASPEr.

Axions and the CMB

Assuming BICEP detected gravitational waves in the CMB (some tension with Planck):

$$H_{\rm inf} \sim 10^{14}\ {\rm GeV}$$

if symmetry broken after inflation → topological defects (strings + domain walls), constrained by observations

if symmetry broken before inflation → inflation can induce isocurvature perturbations of axion, weak constraint on ALPs probed by CASPEr.

for QCD axion, constrains **one** cosmological history.

Requires knowing physics all the way up to GUT scale $\sim 10^{16}\ {\rm GeV}$

many others possible.

QCD Axion and BICEP

Need a high temperature, transient mass, sometime before QCD phase transition.

Need not be on during inflation.



Axion oscillates earlier, damps to high temperature minimum.

Misalignment of minima gives axion dark matter.

Dark matter from choice of parameters instead of initial conditions.

QCD Axion and BICEP

Need a high temperature, transient mass, sometime before QCD phase transition.



e.g. thermal monopole density, Fischler & Preskill (1983)
high temperature mass,
and many others e.g. Kaplan & Zurek (2005), Jeong & Takahashi (2013), G. Dvali (1995)

Bound depends upon high energy physics, while strong CP, axion dark matter rely upon low energy physics.

QCD axion offers unique probe of high energy cosmology,
an era difficult even for gravitational wave detectors

QCD Axion and BICEP

Need a high temperature, transient mass, sometime before QCD phase transition.



e.g. thermal monopole density, Fischler & Preskill (1983)
high temperature mass,
and many others e.g. Kaplan & Zurek (2005), Jeong & Takahashi (2013), G. Dvali (1995)

Bound depends upon high energy physics, while strong CP, axion dark matter rely upon low energy physics.

QCD axion offers unique probe of high energy cosmology,
an era difficult even for gravitational wave detectors

QCD Axion and BICEP

Need a high temperature, transient mass, sometime before QCD phase transition.



e.g. thermal monopole density, Fischler & Preskill (1983)
high temperature mass,
and many others e.g. Kaplan & Zurek (2005), Jeong & Takahashi (2013), G. Dvali (1995)

Bound depends upon high energy physics, while strong CP, axion dark matter rely upon low energy physics.

QCD axion offers unique probe of high energy cosmology,
an era difficult even for gravitational wave detectors